

Layer Preschool




Layer Pre-School provides an environment that is healthy for all attendees (employees & children). In order to maintain such an environment, all staff and children need to be fit, well and in good health; sick children i.e. those who pose a significant risk of infection to others, should remain at home.

Illness or Condition

Exclusion Period

BronchiolitisUntil the child has fully recovered
Chicken pox Until at least 5 days after the appearance of the last crop of spots, when they fully crusted over
ConjunctivitisUntil the eye is treated and appears normal again and after the first course of antibiotics
Diarrhoea and/or vomiting Until 48 Hours from the last episode of diarrhoea or sickness
“Slapped cheek” diseaseUntil the child feels physically well
Hand foot and mouth diseaseUntil all the blisters have healed
Head lice Until the first course of treatment has been given
Hepatitis AUntil the child feels well and for 5 days after the onset of the jaundice
Impetigo Until the infection has completely healed
Measles Until at least 5 days after the rash first appears and the child is well enough
MumpsUntil 5 days after the onset of swollen glands and when child is well enough to return to nursery
RingwormAfter the first course of treatment has been given
Rubella (German measles)Until at least six days after the rash first appears and the child is well enough
Scabies After the first course of treatment has been given
Scarlet fever 48 hours after the first course of antibiotics has been given and the child is well enough
Shingles Exclude if rash is weeping and cannot be covered
Threadworms After the first course of treatment has been given



Layer Preschool



Whooping cough..... 5 days after the first course of antibiotics has been given and the child is well enough

Meningitis viral.....Until the child is well enough

Meningitis (meningococcal)..... At least 48 hours after treatment of antibiotics and only when child is fully recovered.

Meningitis (due to other bacteria)..... At least 48 hours after treatment of antibiotics and only when child is fully recovered.

If a child is found to have an infectious disease during a Pre School session the parent/guardian will be immediately informed and the child will be excluded in accordance with the table above.

If a child is found to have an infectious disease during a Pre-School session the parent/guardian will be immediately informed and the child will be excluded in accordance with the table above.

The Pre-School Child Sickness Procedures

If a child becomes unwell whilst they are at the Pre-School staff are to ensure that:

The parents are contacted immediately

A suitable rest area is allocated for the child

A member of staff remains with the child at all times

Incidents of Vomiting & Diarrhoea Where possible, the Key Person should attempt to ascertain the nature of the child's illness. After the first bout of vomiting, the staff member is to inform the child's parent/guardian and to request that the child is collected from the Pre-School. The parent is asked to keep the child at home for 48 hours after the last bout of vomit or diarrhoea.

High Temperatures Suspicions of high temperature should be checked using an aural thermometer. If the child's temperature is above 38°C the parent/carer is called to advise them of the child's condition.